

An observation on the relation between the fine structure constant and the Gibbs phenomenon in Fourier analysis

Zi-Xiang Zhou

Institute of Mathematics, Fudan University, Shanghai 200433, P.R.China

Email: zxzhou@guomai.sh.cn

Abstract

A value given by a simple mathematical expression is proposed which is close to the fine structure constant given by 1998 CODATA recommended values of the fundamental physical constants up to relative accuracy 10^{-7} . This expression relates closely with the value of the overshoot of the Gibbs phenomenon in Fourier analysis.

1 Introduction

The fine structure constant α is the most important dimensionless universal physical constant. Since it is dimensionless and universal, it is very interesting to know whether it can be expressed by a simple expression of universal mathematical constants. This may help understanding the nature.

As is known, the definition of fine structure constant α is

$$\alpha^{-1} = \frac{2c\epsilon_0 h}{e^2} = \frac{2h}{c\mu_0 e^2} \quad (1)$$

where c is the speed of light in vacuum, μ_0 and $\epsilon_0 = 1/c^2\mu_0$ are the permeability and permittivity of vacuum respectively, e is the elementary charge and h is the Planck constant.

Here we propose

$$\alpha_z^{-1} = \frac{1}{\sqrt{2}} \left(\frac{3\pi}{2} \right)^3 \int_0^\pi \frac{\sin x}{x} dx \approx 137.03598260. \quad (2)$$

This value coincides with the value

$$\alpha^{-1} = 137.03599976 \pm 0.00000050 \quad (3)$$

given by the 1998 CODATA recommended values of the fundamental physical constants [1] up to relative accuracy 1.26×10^{-7} .

In the expression (2), the integral $\text{Si}(\pi) = \int_0^\pi \frac{\sin x}{x} dx$ is related to a universal mathematical constant — the overshoot of the Gibbs phenomenon in Fourier analysis, which is the same to all the jump discontinuities of all piecewisely smooth functions. The expression $\alpha_z^{-1}/\text{Si}(\pi) = (3\pi/2)^3/\sqrt{2}$ is so simple that it seems that there may be physical essence behind the relation $\alpha_z \approx \alpha$, although it is not known presently.

In the next section, the universality of the constant $\text{Si}(\pi)$ in Fourier analysis is reviewed briefly. Then, the value α_z is compared with the values of α given by experiments and physical theories, which were used to determine α for the 1998 CODATA recommended values of the fundamental physical constants. At last, some discussions are presented.

2 A brief review of the Gibbs phenomenon

The Gibbs phenomenon is a universal phenomenon at all the jump discontinuities for Fourier series or Fourier transformations. First, let us look at a simple example. Let $f(x)$ be a square wave of period 2π




Figure 1: Partial sum s_n of the square wave for $n=200$

with $f(x) = 1$ for $0 < x < \pi$ and $f(x) = -1$ for $-\pi < x < 0$. Let s_n be the partial sum of its Fourier series, i.e.

$$s_n(x) = \frac{4}{\pi} \sum_{k=1}^n \frac{\sin(2k-1)x}{2k-1}. \quad (4)$$

Then when $n \rightarrow \infty$, there are overshoots and undershoots in the graph of s_n (Fig. 1).

The limit of the amplitude of s_n near 0 as $n \rightarrow \infty$ is $2C_G$, which is C_G times the jump of $f(x)$ at 0. Here

$$C_G = \frac{2}{\pi} \text{Si}(\pi) = \frac{2}{\pi} \int_0^\pi \frac{\sin x}{x} dx \approx 1.17897974447217. \quad (5)$$

This is the famous Gibbs phenomenon [2, 3, 4, 5], which was discovered more than a century ago.

The Gibbs phenomenon appears not only in the square wave, but also at all jump discontinuities generally. For the Fourier series of any piecewise smooth period function $f(x)$, the limit of the amplitude of the partial sum at a jump discontinuity x_0 of $f(x)$ equals to C_G times the jump of $f(x)$ at that point. That is, for the partial sum $s_n(x)$,

$$\lim_{\delta \rightarrow 0^+} \lim_{n \rightarrow \infty} \text{osc}_{|x-x_0| \leq \delta} s_n(x) = C_G \lim_{\delta \rightarrow 0^+} \text{osc}_{|x-x_0| \leq \delta} \lim_{n \rightarrow \infty} s_n(x) \quad (6)$$

where *osc* refers to the difference of the maximum and the minimum of a function. This fact is also true for the Fourier transformation of a non-periodic function, if the function is absolutely integrable and piecewise smooth [5]. In all cases, the constant C_G is the same.

3 Comparison with experimental values

In Fig. 2, α_z^{-1} given by (2) was compared with the values of α^{-1} given by the experimental data together with various physical theories. Each line segment in the figure represents a datum in Table XV of [1]. That table was used to determine the value α for the 1998 CODATA recommended values of the fundamental physical constants. For each value x with uncertainty σ , the line segment extends from $x - \sigma$ to $x + \sigma$ in Fig. 2. The identifications to the right of the line segments are the same as those in Table XV of [1]. The vertical line marked α_z^{-1} represents the value given by (2) and that marked α^{-1} represents the value given by the 1998 CODATA recommended values of the fundamental physical constants.

4 Discussions

There are other theoretical values of α , such as those given by the string theory [6, 7]. For example, the simplest expression is $\alpha^{-1} = \phi^{-10-\phi^3} = 137.7880938$ where $\phi = (\sqrt{5} - 1)/2$ is the golden mean. More accurate values with accuracy 3×10^{-7} were also given by a complicated expansion [7]. Although the expression α_z^{-1} in this paper is simply an observation, it is much simpler than the other theoretical results with similar accuracy.




Figure 2: Comparison of α_z^{-1} given by (2) with the experimental values which were used to determine α^{-1}

There is a small possibility that $\alpha_z^{-1} \approx \alpha^{-1}$ is simply an accidental coincidence. Actually it is not difficult to construct complicated relations of 2, 3, π etc. to approximate α^{-1} because any real number can be approximated by a rational number to any accuracy. For example, $2^{-19/6} 3^{157/24} \pi^{-1/16} = 137.0360046$. (There is also a number $2^{19/4} 3^{-7/4} 5^{1/4} \pi^{11/4} = 137.036082$ given by Wyler [6].) However, the probability is very small to get accidentally a simple relation like $\alpha_z^{-1}/\text{Si}(\pi) = (3\pi/2)^3/\sqrt{2}$ in this paper. The internal relation between α_z and the fine structure constant is to be revealed.

References

- [1] P. J. Mohr and B. N. Taylor, Rev. Mod. Phys. **72** (2000) 351 ; J. Phys. & Chem. Ref. Data **28** (1999) 1713.
- [2] E. Hewitt and R. E. Hewitt, Arch. Hist. Exact Sci. **21** (1979) 129.
- [3] T. W. Körner, *Fourier analysis* (Cambridge University Press, Cambridge, 1988).
- [4] J. S. Walker, *Fourier analysis* (Oxford University Press, Oxford, 1988).
- [5] D. C. Champeney, *A handbook of Fourier theorems* (Cambridge University Press, Cambridge, 1987).
- [6] M. S. El Naschie, Chaos, Solitons and Fractals **10** (1999) 1947.
- [7] M. S. El Naschie, Chaos, Solitons and Fractals **12** (2001) 801.